



## Wolcott Rallison Syndrome caused by a Novel Mutation in EIF2AK3 Gene

Vikram Bhaskar<sup>\*1</sup>, Chennakeshava Thunga<sup>1</sup>, Soumya Tiwari<sup>2</sup>, Sarah Flanagan<sup>3</sup>, Sean Ellard<sup>3</sup>

<sup>1</sup>(M.D. pediatrics), Senior resident, Lady Hardinge Medical College, India

<sup>2</sup>Associate Professor, Lady Hardinge Medical College, India

<sup>3</sup>Royal Devon and Exeter NHS Healthcare Trust, Exeter, UK

### Abstract

Wolcott-Rallison syndrome (WRS) is an autosomal recessive disorder characterized by early-onset diabetes, skeletal dysplasia, and growth retardation<sup>1</sup>. Some patients of WRS may develop central hypothyroidism, hepatic dysfunction, renal insufficiency or central nervous system abnormalities<sup>2,3</sup>. This is a very rare disorder and, less than 100 cases have been described till 2009<sup>4</sup>. Mutations in the eukaryotic translation initiation factor 2 $\alpha$  kinase (EIF2AK3) gene are responsible for this disorders<sup>5</sup>. We hereby report a case of Wolcott-Rallison syndrome caused by a novel mutation in EIF2AK3 gene

**Keywords:** Wolcott rallison, Neonatal diabetes, Beta cell disorders

### Corresponding author: Vikram Bhaskar

(M.D. pediatrics), Senior resident, Lady Hardinge Medical College, India. Email: [vicky\\_8884@yahoo.co.in](mailto:vicky_8884@yahoo.co.in)

**Citation:** Vikram Bhaskar et al. (2018), Wolcott Rallison Syndrome caused by a Novel Mutation in EIF2AK3 gene. Int J Ped & Neo Heal. 2:4, 52-53

**Copyright:** ©2018 Vikram Bhaskar et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

**Received:** March 07, 2018

**Accepted:** March 17, 2018

**Published:** April 30, 2018

### Introduction

A 5-month-old female child was brought to our hospital with complaints of multiple episodes of multi-focal tonic clonic seizures for the last 1 day. She was the first in order of birth, born from a non-consanguineous marriage, delivered at term with weight appropriate for gestational age, without significant perinatal problems. There was no significant family history.

On routine investigations patient was found to have persistent hyperglycemia. Other investigations including CSF examination, liver and kidney function tests, skull ultrasound and serum electrolytes were within normal limit. All cultures including blood, CSF, and urine culture were sterile. We investigated her further for hyperglycemia and found that her C-peptide levels were inappropriately low (<0.30 ng/mL) for her blood sugar levels. Considering the possibility of infantile diabetes, her blood samples were sent to Royal Devon and Exeter NHS foundation trust, Exeter, UK, for genetic testing. Analysis of exons 1, 6, 7, 9, 11,

14, 16 and 17 of the EIF2AK3 gene (AF110146.1) was done by Sanger sequencing as previously described (Rubio-Cabezas JCEM 2009). Primer sequences are available on request. She was found to be homozygous for a novel EIF2AK3 missense mutation, p.R1064Q (c.3191G>A), in exon 17. This mutation has never been reported before, but it affects a highly conserved nucleotide and in silico evidence suggest that it is likely to be pathogenic. This result is consistent with a diagnosis of Wolcott Rallison syndrome. Both the parents were found to be heterozygous for this novel mutation.

The infant was managed initially with intravenous regular insulin infusion, and later shifted to subcutaneous NPH insulin at discharge. She remains under follow up, with reasonable glycemic control (on NPH insulin @ 0.5 IU/kg/day), and no episode of ketoacidosis or hypoglycemia. Currently she is 13 months old, and till date there is no evidence of skeletal dysplasia or hepatic and renal dysfunction. Mild motor delay was present in development. Seizures are controlled on oral phenytoin.

### Discussion:

Mutations in the gene encoding the EIF2AK3 protein are responsible for Wolcott Rallison syndrome. Our patient was found to be homozygous for a novel missense mutation (p.R1064Q) in exon 17 of the EIF2AK3 gene. Her mother and father were confirmed to be heterozygous carriers.

The age at onset of diabetes in Wolcott-Rallison syndrome is variable, ranging from 2 weeks to 30 months in some cases<sup>6,7</sup>. Our patient presented at the age of 5 months. Early onset, insulin dependent diabetes is present in all the patients of this syndrome<sup>8</sup>.

Clinically, our patient presented with history of seizures, and had developmental delay (motor), however there was no evidence of any liver, kidney or skeletal involvement till date, but these manifestations may be delayed by several years<sup>8</sup>. Developmental delay is a common finding in WRS, and its severity is variable. Severe cases have been reported with neuro-motor deficit and epilepsy<sup>9</sup>.

Making a genetic diagnosis in patients with WRS is important not only for genetic counseling, but also because the disease has a multisystem involvement and variable presentation, and thus patients can be monitored for various system complications.

**References:**

1. Wolcott CD, Rallison ML. [Infancy-onset diabetes mellitus and multiple epiphyseal dysplasia](#). *J Pediatr*.1972;80:292–297.
2. Bin-Abbas B, Al-Mulhim A, Al-Ashwal A. [Wolcott-Rallison syndrome in two siblings with isolated central hypothyroidism](#). *Am J Med Genet*. 2002;111:187–190.
3. Engelmann G, Meyburg J, Shahbek N, Al-Ali M, Hairetis MH, Baker AJ, Rodenburg RJT, Wenning D, Flechtenmacher C, Ellard S, Smeitink JA, Hoffmann GF, Buchanan CR. [Recurrent acute liver failure and mitochondriopathy in a case of Wolcott-Rallison syndrome](#). *J Inherit Metab Dis*. 2008;31:540–546.
4. Rubio-Cabezas O, Patch AM, Minton JA, Flanagan SE, Edghill EL, Hussain K, Balafrej A, Deeb A, Buchanan CR, Jefferson IG, Hattersley AT, Ellard S. [Wolcott-Rallison syndrome is the most common genetic cause of permanent neonatal diabetes in consanguineous families](#). *J Clin Endocrinol Metab*.2009;94:4162–4170.
5. Delepine M, Nicolino M, Golamaully M, Lathrop GM, Julier C. [EIF2AK3, encoding translation initiation factor 2-alpha kinase 3, is mutated in patients with Wolcott-Rallison syndrome](#). *Nat Genet*.2000;25:406–409.
6. Castelnau P, Le Merrer M, Diatloff-Zito C, Marquis E, Tête M-J, Robert J-J. [Wolcott–Rallison syndrome: A case with endocrine and exocrine pancreatic deficiency and pancreatic hypotrophy](#). *Eur J Pediatr*. 2000;159:631–633.
7. Ozbek MN, Senée V, Aydemir S, Kotan LD, Mungan NO, Yuksel B, Julier C, Topaloglu AK. [Wolcott–Rallison syndrome due to the same mutation \(W522X\) in EIF2AK3 in two unrelated families and review of the literature](#). *Pediatr Diabetes*. 2010;11:279–285.
8. Iyer S, Korada M, Rainbow L, Kirk J, Brown RM, Shaw N, Barrett TG. [Wolcott-Rallison syndrome: a clinical and genetic study of three children, novel mutation in EIF2AK3 and review of the literature](#). *Acta Paediatr*. 2004;93:1195–1201.
9. De Wit MC, de Coo IF, Julier C, Delepine M, Lequin MH, van de Laar I, Sibbles BJ, Bruining GJ, Mancini GM: [Microcephaly and simplified gyral pattern of the brain associated with early onset insulin-dependent diabetes mellitus](#). *Neurogenetics* 2006, 7:259–263.